

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 092 776 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

18.04.2001 Bulletin 2001/16

(21) Application number: 00121763.7

(22) Date of filing: 05.10.2000

(51) Int. Cl.⁷: **C12N 15/52**, C12N 9/00,
C12P 13/06, C12P 13/08,
C12P 13/14, C12R 1/19
// (C12P13/06, C12R1:19),
(C12P13/08, C12R1:19),
(C12P13/14, C12R1:19)

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

Designated Extension States:

AL LT LV MK RO SI

(30) Priority: 14.10.1999 RU 99121636

(71) Applicant: Ajinomoto Co., Inc.

Tokyo (JP)

(72) Inventors:

• Gusyatiner, Mikhail Marovich

113535 Moscow (RU)

• Kozlov, Yuri Ivanovich

117574 Moscow (RU)

• Ptitsyn, Leonid Romanovich

115404 Moscow (RU)

• Altman, Irina Borisovna

119435 Moscow (RU)

• Voroshilova, Elvira Borisovna

113535 Moscow (RU)

• Iomantas, Yurgis Antanas Vladovich

127273 Moscow (RU)

• Yampolskaya, Tatyana Abramovna

123364 Moscow (RU)

(74) Representative:

Strehl Schübel-Hopf & Partner

Maximilianstrasse 54

80538 München (DE)

(54) **Method for producing L-amino acid by fermentation**

(57) L-amino acid such as L-threonine and L-glutamic acid is produced by cultivating a bacterium belonging to the genus *Escherichia* which is transformed with a gene coding for pyruvate carboxylase and has L-amino acid productivity; producing and accumulating L-amino acid in the medium and collecting the L-amino acid from the medium.

EP 1 092 776 A1

Description

Technical Field

5 [0001] The present invention relates to a method for producing L-amino acid, specifically relates to a method for producing L-threonine, L-glutamic acid, L-homoserine, L-methionine, L-arginine, L-proline or L-isoleucine using a bacterium belonging to the genus *Escherichia*.

Background Art

10 [0002] L-amino acids such as L-threonine and L-glutamic acid have conventionally been produced by fermentation methods principally utilizing coryneform bacteria which belong to the genera *Brevibacterium*, *Corynebacterium*, and *Microbacterium* or mutants thereof ("Amino Acid Fermentation", Gakkai Shuppan Center, pp.195-215, 1986).

[0003] On the other hand, there has been disclosed techniques for breeding of L-amino acid-producing bacteria 15 belonging to the genus *Escherichia* utilizing genetic recombination techniques.

[0004] For example, there has been disclosed a method of producing L-lysine using *Escherichia coli* in which genes coding for dihydrodipicolinate synthetase, aspartokinase are desensitized to feedback inhibition by L-lysine and L-threonine, and diaminopimelate dehydrogenase (or tetrahydrodipicolinate succinylase and succinyldiaminopimelate 20 deacylase) are enhanced (WO 95/16042).

[0005] Though the productivity of L-amino acids has considerably been improved by breeding of such microorganisms as mentioned above or production processes have been improved, it is still desired to develop more efficient processes for producing L-amino acids in order to meet the expected markedly increased future demand of the amino acids.

[0006] Pyruvate carboxylase (pyruvate:carbon dioxide ligase (ADP-forming), EC 6.4.1.1] is a biotine containing enzyme. It catalyses the carboxylation of pyruvate to form oxaloacetate. Pyruvate carboxylase has not been known in 25 *Escherichia coli*, but *Bacillus subtilis* (Diesterhaft, M.D. and Freese, E., J. Biol. Chem., 248, No.17, 6062-6070 (1973)) and *Corynebacterium glutamicum* (Peters-Wendisch, P. G. et al., *Microbiology*, 144, 915-927 (1998)) have pyruvate carboxylase and the nucleotide sequences of the genes coding for the enzymes have been reported.

[0007] Further, there has been known *Corynebacterium glutamicum* in which an activity of pyruvate carboxylase is enhanced or *Corynebacterium glutamicum* of which pyruvate carboxylase gene is inactivated (Peters-Wendisch, P. G. 30 et al., *Microbiology*, 144, 915-927 (1998)). However, these microorganisms were created as experimental materials to study enzymes necessary for growth with glucose. The relation between the activity of pyruvate carboxylase and productivity of L-amino acids has not been known. Also, there has not been known an attempt to introduce the pyruvate carboxylase gene into a bacterium belonging to the genus *Escherichia*.

Summary of the Invention

[0008] The present invention has been accomplished from the aforementioned point of view, and its object is to provide a method for producing L-amino acid, especially L-threonine, L-lysine, L-glutamic acid, L-homoserine, L-methionine, L-arginine, L-proline or L-isoleucine in high efficiency.

40 [0009] As a result of diligent investigation in order to achieve the object described above, the present inventors have found that introduction of a gene coding for pyruvate carboxylase (hereinafter referred to as "pyc gene") into a bacterium belonging to the genus *Escherichia* can improve L-amino acid productivity of the bacterium. Thus, the present invention has been completed.

[0010] That is, the present invention is a bacterium belonging to the genus *Escherichia* which is transformed with 45 a gene coding for pyruvate carboxylase and has L-amino acid productivity.

[0011] The present invention also provides the bacterium described above wherein the gene coding for pyruvate carboxylase is derived from a bacterium belonging to the genus *Bacillus*.

[0012] The present invention further provides the bacterium described above wherein the gene coding for pyruvate carboxylase is introduced into said bacterium in a low copy number.

50 [0013] The present invention still further provides the bacterium described above wherein the L-amino acid is selected from the group consisting of L-threonine, L-lysine, L-glutamic acid, L-homoserine, L-methionine, L-arginine, L-proline and L-isoleucine.

[0014] The present invention also provides a method for producing L-amino acid comprising cultivating the above bacterium in a medium, producing and accumulating the L-amino acid in the medium, and collecting the L-amino acid 55 from the medium.

[0015] The present invention further provides the method described above wherein the L-amino acid is selected the group consisting of L-threonine, L-lysine, L-glutamic acid, L-homoserine, L-methionine, L-arginine, L-proline and L-isoleucine.

[0016] The term "L-amino acid productivity" means a property that a bacterium produces and accumulates the L-amino acid in a medium in a larger amount than a wild type strain thereof.

[0017] According to the present invention, productivity of L-amino acid such as L-threonine, L-lysine, L-glutamic acid,

5 L-homoserine, L-methionine, L-arginine, L-proline or L-isoleucine can be improved in a bacterium belonging to the genus *Escherichia*. Further, the present invention may be utilized to breeding of an L-amino acid producer belonging to the genus *Escherichia*.

Brief Description of the Drawings

[0018]

Figure 1 shows a scheme of constructing *Bacillus subtilis* pycA::KmR recipient strain for cloning pycA gene,

Figure 2 shows a scheme of cloning pycA gene from *Bacillus subtilis* 168, and

15 Figure 3 shows a scheme of construction of the plasmid containing pycA gene.

Detailed Description of the Invention

[0019] The present invention will be explained in detail below.

(1) A bacterium belonging to the genus *Escherichia* transformed with pyc gene

[0020] A bacterium belonging to the genus *Escherichia* of the present invention is a bacterium transformed with pyc gene. As the bacterium belonging to the genus *Escherichia*, there exemplified by *Escherichia coli*. While the pyc gene used in the present invention is not particularly limited so long as it can code a protein having pyruvate carboxylase activity, examples of the pyc gene include, for example, pyc gene derived from bacteria belonging to the genus *Bacillus* or pyc gene from coryneform bacteria. The nucleotide sequences of the pyc gene derived from *Bacillus subtilis* (Genbank/EMBL/DDBJ Accession Z97025, NID g2224758) and *Corynebacterium glutamicum* (Peters-Wendisch, P. G. et al., Microbiology, 144, 915-927 (1998)) has been reported. Therefore, pyc gene can be obtained by PCR (polymerase chain reaction: White, T.J. et al., Trends Genet., 5, 185 (1989)) using primers synthesized according to the nucleotide sequences from chromosomal DNA of bacteria belonging to the genus *Bacillus* such as *Bacillus subtilis* or coryneform bacteria such as *Corynebacterium glutamicum* as a template.

[0021] A transformant of *Escherichia coli* 44/pMW119-pycA, which harbors pMW119-pycA containing pyc gene of *Bacillus subtilis* has been deposited in Russian National Collection of Industrial Microorganisms (VKPM) Depository, GNIIGenetika; 1, Dorozhny Proezd., 1, 113545, Moscow, Russia, since August 30, 1999 under a Registration number of VKPM B-7822, and transferred from the original deposit to international deposit based on Budapest Treaty on September 1, 2000.

[0022] A transformant of *Escherichia coli* MG442/pMW119-pycA, which harbors pMW119-pycA containing pyc gene of *Bacillus subtilis* has been deposited in Russian National Collection of Industrial Microorganisms (VKPM) Depository, GNIIGenetika; 1, Dorozhny Proezd., 1, 113545, Moscow, Russia, since August 30, 1999 under a Registration number of VKPM B-7821, and transferred from the original deposit to international deposit based on Budapest Treaty on September 1, 2000.

[0023] Incidentally, the gene coding pyruvate carboxylase of *Bacillus subtilis* (pycA gene) has been cloned as described below in a process of accomplishing the present invention.

[0024] First, a pycA gene-deficient strain of *Bacillus subtilis* was constructed. Then, using the strain as a recipient, pycA gene was cloned by isolating a DNA fragment which complemented citrate or aspartate auxotrophy of the strain from a genomic DNA library of a wild type strain of *Bacillus subtilis*. Since *Bacillus subtilis* forms oxaloacetate by pyruvate carboxylase, *Bacillus subtilis* cannot grow on minimal medium if it is deficient in the enzyme. However the growth is repaired by addition of L-aspartate, L-glutamate, citrate or succinate. The nucleotide sequence of pycA gene and the amino acid sequence coded by the gene are shown in SEQ ID Nos: 3 and 4.

[0025] The pycA gene-deficient strain is obtained, for example, as described below. A partial DNA fragment of ylaP gene located immediately after the pycA gene on the chromosome is obtained by PCR. As primers for PCR are exemplified by oligonucleotides of SEQ ID NOs: 1 and 2. Then a wild type strain of *Bacillus subtilis* is transformed with a plasmid DNA containing the obtained DNA fragment and a marker gene, followed by integrating the plasmid DNA into ylaP gene on the chromosomal DNA by homologous recombination. Next, chromosomal DNA library of the obtained strain is constructed using a restriction enzyme which recognizes a restriction site inside of pycA gene, for example, *Pst*I. The recombinant plasmid containing the marker gene and a partial pycA gene fragment may be obtained by transforming a wild type strain of *Bacillus subtilis* with the library and selecting clones expressing the marker gene. Then a wild type

strain of *Bacillus subtilis* is transformed with the linearized recombinant plasmid and clones expressing the marker gene are selected to obtain *pycA* gene-deficient strain in which the plasmid DNA is integrated in *pycA* gene on chromosomal DNA of the strain.

[0026] Transformation of a bacterium belonging to the genus *Escherichia* with *pyc* gene may be performed by inserting *pyc* gene into an appropriate vector which functions in a bacterium belonging to the genus *Escherichia* to construct a recombinant vector, and introducing the recombinant vector to a bacterium belonging to the genus *Escherichia*.

[0027] The vector is exemplified by plasmid such as pBR322, pMW118, pMW119, pUC18, pUC19 or the like, phage vector such as λ 1059, λ BF101, M13mp9 or the like, and transposon such as Mu, Tn10, Tn5 or the like. However, in the present invention, a low copy plasmid such as pMW118 and pMW119 is preferable, because a transformant harboring a vector may be unstable if a high copy plasmid is used to introduce *pycA* gene as a vector.

[0028] The introduction of a DNA into a bacterium belonging to the genus *Escherichia* can be performed, for example, by a method of D. A. Morrison (Methods in Enzymology 68, 326 (1979)) or a method in which recipient bacterial cells are treated with calcium chloride to increase permeability of DNA (Mandel, M. and Higa, A., J. Mol. Biol., 53, 159 (1970)) and the like.

[0029] Preparation of the genomic DNA, construction of genomic DNA library, hybridization, PCR, preparation of plasmid DNA, digestion and ligation of DNA, transformation and the like are performed according to the methods which are known to a person skilled in the art. Such methods were described by Sambrook, J., Fritsche, E. F., Maniatis, T. in Molecular Cloning, Cold Spring Harbor Laboratory Press, 1.21 (1989).

[0030] L-amino acid productivity of a bacterium belonging to the genus *Escherichia* can be improved by conferring or enhancing pyruvate carboxylase activity on the bacterium through transforming the bacterium with *pyc* gene. This is probably because oxaloacetic acid is formed not only from phosphoenolpyruvic acid but also from pyruvic acid. That is, transport of glucose (or sucrose) molecule to a bacterial cell which is mediated by phosphoenolpyruvic acid system proceeds by consuming one molecule of phosphoenolpyruvic acid being accompanied with formation of one molecule of pyruvic acid. Although formed pyruvic acid is not directly utilized for L-amino acid synthesis in a bacterial cell, L-amino acid productivity may be increased by formation of oxaloacetic acid from pyruvic acid.

[0031] As a bacterium belonging to the genus *Escherichia* to which *pyc* gene is to be introduced, a strain having an ability to produce objective L-amino acid is used. Alternatively, a bacterium belonging to the genus *Escherichia* which is transformed with *pyc* gene may be conferred L-amino acid productivity. Incidentally, *pyc* gene of *Escherichia coli* has not been known, however, if the gene is found in *E. coli* hereafter, the gene may be used.

[0032] Examples of L-amino acid-producing bacteria belonging to the genus *Escherichia* are described below.

(1) L-threonine-producing bacteria

[0033] The L-threonine-producing bacteria belonging to the genus *Escherichia* may be exemplified by strain MG442 (deposited in Russian National Collection of Industrial Microorganisms (VKPM) Depositary, GNIIGenetika; 1, Dorozhny Proezd., 1, 113545, Moscow, Russia, under a Registration number of VKPM B-1628)(USP 4,278,765; Guayatinier M.M. et al., *Genetika* (in Russian), 14, 947-956 (1978)). The strain MG442 had been obtained as a mutant resistant to L-threonine analogue, 2-amino-3-hydroxyvaleric acid, it was also an isoleucine auxotroph of leaky type. It produces L-threonine and L-glutamate as a by-product.

[0034] Alternatively, a transformant of MG442 which is transformed with pVIC40 (USP 5,175,107) may be preferably used as L-threonine-producing strain.

(2) L-Lysine-producing bacteria

[0035] The L-lysine-producing bacteria belonging to the genus *Escherichia* may be exemplified by various bacterium described in WO95/16042. The L-lysine-producing bacteria are concretely exemplified by a bacterium in which aspartokinase activity and dihydrodipicolinate reductase activity are increased such as W3110(*tyrA*)RSFD80+pdapB.

(3) L-homoserine-producing bacteria

[0036] The L-homoserine producing bacteria belonging to the genus *Escherichia* may be exemplified by strain 44 (*thrB*). This strain was derived from the known strain C600 (*thrB*, *leuB*) (Appleyard R.K., Genetics, 39, 440-452 (1954)) as *Leu*⁺ revertant. A transformant strain of 44 being transformed with a plasmid containing a *thrA* gene coding for aspartokinase-homoserine dehydrogenase I may be preferably used.

[0037] A strain of *Escherichia coli* 44 has been deposited in Russian National Collection of Industrial Microorganisms (VKPM) Depositary, GNIIGenetika; 1, Dorozhny Proezd., 1, 113545, Moscow, Russia, since September 23, 1980 under a Registration number of VKPM B-2175, and transferred from the original deposit to international deposit based on Budapest Treaty on September 1, 2000.

(4) L-glutamic acid-producing bacteria

[0038] The L-glutamic acid producing bacteria belonging to the genus *Escherichia* may be exemplified by a strain which is resistant to L-valine, for example, *Escherichia coli* strains B11, K-12 (ATCC10798), B (ATCC11303) and W (ATCC9637).

(5) L-isoleucine-producing bacteria

[0039] The L-isoleucine-producing bacteria belonging to the genus *Escherichia* may be exemplified by a strain TDH6/pVIC40, pMWD5 (Hashiguchi, K. et al., Biosci. Biotechnol. Biochem., 63(4), 672-679 (1999)) or AJ12919 described in EP0 685 555 A1.

(2) Method for producing an L-amino acid

[0040] An L-amino acid can be efficiently produced by cultivating a bacterium belonging to the genus *Escherichia* which is transformed with *pycA* gene as described above and has an L-amino acid productivity in an appropriate medium, producing and accumulating the L-amino acid in the medium, and collecting the L-amino acid from the medium.

[0041] The L-amino acid is exemplified preferably by L-threonine, L-lysine, L-glutamic acid, L-homoserine, L-methionine, L-arginine, L-proline and L-isoleucine. The L-amino acid may be produced by the method of the present invention either alone or as a mixture of more than two kinds of L-amino acids.

[0042] In the method of present invention, the cultivation of the bacterium belonging to the genus *Escherichia*, the collection and purification of amino acid from the liquid medium may be performed in a manner similar to those of the conventional method for producing an amino acid by fermentation using a bacterium. A medium used in cultivation may be either a synthetic medium or a natural medium, so long as the medium includes a carbon and a nitrogen source and minerals and, if necessary, nutrients which the bacterium used requires for growth in appropriate amounts. The carbon source may include various carbohydrates such as glucose and sucrose, and various organic acids. Depending on assimilatory ability of the used bacterium, alcohol including ethanol and glycerol may be used. As the nitrogen source, ammonia, various ammonium salts such as ammonium sulfate, other nitrogen compounds such as amines, a natural nitrogen source such as peptone, soybean hydrolyzate and digested fermentative microbe are used. As minerals, monopotassium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, calcium carbonate are used.

[0043] The cultivation is preferably performed under an aerobic condition such as a shaking culture, and an aeration and stirring culture. The temperature of culture is usually 20 to 40°C, preferably 30 to 38°C. The pH of the culture is usually between 5 and 9, preferably between 6.5 and 7.2. The pH of the culture can be adjusted with ammonia, calcium carbonate, various acids, various bases, and buffers. Usually, 1 to 3-day cultivation leads to the accumulation of the target L-amino acid in the medium.

[0044] Collecting and purifying the L-amino acid can be performed by removing solids such as cells from the medium by centrifugation or membrane filtration after cultivation, followed by ion exchange, concentration and crystalline fraction methods and the like.

Best Mode for Carrying Out the Invention

[0045] The present invention will be more concretely explained below with reference to Examples.

Example 1: Cloning of the *pycA* gene from *Bacillus subtilis* 168 strain(1) Construction of *B. subtilis* *pycA*::KmR recipient strain for cloning

[0046] The recipient strain of *B. subtilis* was constructed by insertional mutagenesis of *pycA* gene. The fragment of *pycA* gene was cloned in two steps (refer to Figs. 1 and 2).

[0047] In a data bank one of reading frame (ylaP) from *B. subtilis* genome sequence project is identified as *pycA* gene, encoding pyruvate carboxylase. The region located immediately after the *pycA* gene terminator was cloned by PCR amplification, using the following pair of oligonucleotides:

5' -GATATAAGGGGACTTCAGAG, and
5' -GGCGCTTTATGCGTTTCAATC.

[0048] The 269bp DNA fragment was blunt-ended with Klenow fragment of DNA polymerase I and cloned into *ScaI* site of pBR322 plasmid in *E. coli* strain. The resulting plasmid pBRPYCA11 was digested by *ClaI* and ligated to 1628bp DNA fragment carrying *cat*-gene, which was obtained by digestion of pC194 with *ClaI*. Then, using the obtained plasmid pBRPYCA11CmR3, *Bacillus subtilis* 168 trpC2 strain was transformed and Cm^R-clones were selected. The plasmid pBRPYCA11CmR3 cannot autonomously replicate in bacterium belonging to the genus *Bacillus*, but can integrate into chromosome near the *pycA* gene by homology with the 269bp DNA fragment. The strain was named as *B. subtilis* trpC2 xyz::pBRPYCACmR3. The integrant carried pBR322-replicon, *tet* gene encoding resistance to tetracycline derived from pBR322 and Cm^R marker.

[0049] Chromosomal DNA of *B. subtilis* trpC2 xyz::pBRPYCACmR3 was isolated and used to clone a distal part of *pycA* gene. The chromosomal DNA which is digested with *PstI* was self-ligated and transformed into *E. coli* TG1 strain. Recombinants resistant to tetracycline and chloramphenicol were selected. The plasmid was recovered from one of the recombinant and named as pPYC1. The plasmid pPYC1 contained a single *BglII* restriction site, which located in *pycA* gene fragment. This site was used to insert 1818bp DNA fragment, containing a kanamycin resistance marker for *B. subtilis*. The plasmid pPYC1 was digested with *BglII* and ligated to pUC7KmR which was digested with *BamHI* to obtain a plasmid pPYC1::KmR.

[0050] The inactivation of *pycA* gene in a wild type *B. subtilis* 168 trpC2 was achieved by transformation of the same strain with linearized plasmid pPYC1::KmR by one step procedure, followed by selection on Luria broth supplemented with kanamycin. The Km^R marker was inserted into *pycA* gene on chromosomal DNA by homologous double-crossover recombination.

[0051] The constructed strain of *B. subtilis* *pycA*::Km^R trpC2 did not grow in Spizizen minimal medium (Anagnostopoulos, C and Spizizen, J., J. Bacteriol., 81, 741-746 (1961)) with tryptophan unless citrate or aspartate were added to the media.

(Composition of Spizizen minimal medium)

[0052]

K ₂ HPO ₄ • 3H ₂ O	18.3 g/L
KH ₂ PO ₄	6.0 g/L
(NH ₄) ₂ SO ₄	2.0 g/L
sodium citrate • 5H ₂ O	1.2 g/L
MgSO ₄ • 7H ₂ O	0.2 g/L
Glucose	10.0 g/L
pH7.0	

[0053] The *B. subtilis* trpC2 *pycA*::Km^R *recE*::Tc^R strain was constructed by transducing the inactivated allele of *recE* gene from *B. subtilis* *recE*::Tc^R strain with phage E40 (Fig. 2). It was necessary to inactivate recombination in recipient strain for cloning of *pycA* gene from *B. subtilis*. The constructed strain *B. subtilis* trpC2 *pycA*::Km^R *recE*::Tc^R strain was used as recipient to clone *pycA* gene.

(2) Cloning of *B. subtilis* native *pycA* gene by complementation.

[0054] The *pycA* gene was cloned from *B. subtilis* 168 trpC2 strain. Chromosomal DNA prepared from the strain 168 trpC2 was partially digested with *EcoRI* and ligated to pCB20 (Em^R) (gifted by Dr. Yu Jomantas, refer to "Genetic transformation and expression" L.O. Butler, et al. eds., Intercept Ltd., PO Box 402, Wimborne, Dorset, BH229T2, U.K., 1989, p269-281) which was digested with *EcoRI*. Then the trpC2 *pycA*::Km^R *recE*::Tc^R recipient strain was transformed with the ligation mixture (Fig. 2). The Asp⁺, Em^R, Km^R clones were selected on Spizizen minimal medium without citrate or aspartate supplementation but supplemented with tryptophan, erythromycin and kanamycin. The plasmid pPYCR3 harbored by one of the selected clones complemented *pycA* mutation, had a predicted structure and conferred the same Asp⁺, Em^R phenotype after second transformation.

(3) Cloning of DNA fragment carrying *pycA* gene from *B. subtilis* 168 trpC2 strain into high copy number plasmid pUC18 and pUC19

[0055] Plasmid pPYCR3 was digested with *Hind*II and *Sca*I and *Hind*II-*Sca*I fragment (4414 bp), carrying *pycA* gene with 5'-end upstream sequence (including promoter and ribosome binding site), was cloned in pUC18 digested with *Sma*I. Thus the plasmid pUC18-*pycA* was obtained (Fig. 3, upside). The ligation reaction was performed concretely as follows. Ninety ng of *Sma*I-digested pUC18 and 300 ng of *Hind*II-*Sca*I fragment (4414 bp) carrying *pycA* gene which was purified by agarose gel electrophoresis were ligated with 2 units of T4 DNA Ligase (Pharmacia, Sweden) in 45 µl of reaction mixture containing 1x One-Phor-All buffer (Pharmacia, Sweden) and 1 mM ATP.

[0056] *E. coli* TG1 was transformed with pUC18-*pycA*. Obtained transformants were analyzed by using *Kpn*I, *Xba*I, *Bam*HI, *Hind*III and *Eco*RI endonucleases. All clones carried the *pycA* gene in opposite direction to *lac* promoter of pUC18 plasmid. However, the *pycA* genes cloned in pUC18 may be expressed under the control of the inherent promoter of the gene.

[0057] Then in order to rearrange *pycA* gene under control of *lac* promoter, the *Kpn*I-*Xba*I fragment of pUC18-*pycA* was cloned into pUC19 digested with *Kpn*I-*Xba*I. However, the obtained clones were unstable and the recombinant plasmid was eliminated during short time (6-8 h) of cultivation.

(4) Cloning of DNA fragment carrying *pycA* gene into a low copy number plasmid pMW119

[0058] The *Kpn*I-*Xba*I fragment of pUC18-*pycA* was ligated to a low copy number plasmid pMW119 digested with *Kpn*I and *Xba*I (Fig. 3, downside). The ligation reaction was performed in 50 µl of reaction mixture containing 60 ng of pMW119 digested with *Kpn*I-*Xba*I, 120 ng of *Kpn*I-*Xba*I fragment of pUC18-*pycA* which was purified by agarose gel electrophoresis, 1x One-Phor-All buffer (Pharmacia, Sweden), 1 mM ATP and 1 unit of T4 DNA Ligase (Pharmacia, Sweden).

[0059] Using the ligation mixture, *E. coli* TG1 was transformed. Resulting transformants were analyzed using *Sac*I, *Kpn*I, *Xba*I, *Bam*HI, *Hind*III and *Eco*RI. All clones carried *pycA* gene under control of *lac* promoter of pMW119 and its own promoter. The thus obtained clone was named TG1(pMW119-*pycA*), from which the plasmid pMW119-*pycA* was obtained.

Example 2: Production of L-threonine and L-glutamic acid

[0060] *E. coli* strain MG442 was transformed with the plasmid pMW119-*pycA* and 10 Amp^r colonies were selected. Productivities of L-threonine and L-glutamic acid of the colonies were examined using the following fermentation medium.

(Composition of fermentation medium)

[0061]

glucose (separately sterilized)	60 g/L
(NH ₄) ₂ SO ₄	15.0 g/L
KH ₂ PO ₄	1.5 g/L
MgSO ₄ · 7H ₂ O	1.0 g/L
L-isoleucine	100 mg/L
thiamine	0.1 mg/L
CaCO ₃ (separately sterilized)	20 g/L

[0062] Fermentation was performed in test tubes (20x300mm) at 32 °C for 72 hours using a rotary shaker. One loop of each of 10 clones was inoculated into the test tube containing 2.0 ml of the fermentation medium. As controls, 10 single colonies of a plasmid-less strain MG442 were used. After the fermentation, concentrations of L-threonine and L-glutamic acid accumulated in the media were measured by thin-layer chromatography. The results are shown in Table 1.

Table 1

Clone Nos.	MG442(control)		MG442(pMW119-pycA)	
	L-thr G/L	L-glu G/L	L-thr g/L	L-glu g/L
1	5.5	9.5	7.8	12.3
2	3.2	7.1	9.8	12.3
3	3.6	7.1	9.4	13.3
4	3.4	8.9	9.8	11.8
5	3.3	8.9	9.8	11.8
6	4.8	7.3	9.9	11.8
7	3.3	7.0	10.0	10.7
8	4.7	8.3	10.0	10.4
9	3.2	7.3	9.7	11.2
10	2.6	4.1	10.0	10.7
Average	3.8	7.4	9.6	11.6

Example 3: Production of L-homoserine

[0063] L-homoserine-producing E. coli strain 44 was transformed with the plasmid pMW119-pycA and five ampiciline resistant clones were selected. Production of L-homoserine was examined as it is indicated in the Example 1, but the fermentation medium was supplemented with 0.5 g/l of L-threonine. As a control five colonies of plasmidless strain 44 were used. After the fermentation average L-homoserine concentrations were 4.5 g/l (the strain 44) and 5.7 g/l (the strain 44 with the plasmid pMW119-pycA).

[0064] Also the effect of pycA gene on the productivity of L-lysine is reasonably expected, because pycA gene is confirmed to have the positive effect on the productivity of L-threonine and L-homoserine which share the same biosynthetic pathway in part as L-lysine.

SEQUENCE LISTING

5 <110> Ajinomoto Co., Inc.
 <120> Method for Producing L- Amino Acid by Fermentation
 10 <130> EPA-53466

15 <150> RU 99121636
 20 <151> 1999-10-14

<160> 4
 25 <170> Patentln Ver. 2.0

30 <210> 1
 <211> 20
 <212> DNA
 35 <213> Artificial Sequence

<220>
 40 <223> Description of Artificial Sequence:primer

<400> 1
 gatataaggg gacttcagag 20
 45

<210> 2
 <211> 20
 50 <212> DNA
 <213> Artificial Sequence

55

<220>

<223> Description of Artificial Sequence:primer

<400> 2

ggcgctttat gcgtttcaatc

20

<210> 3

<211> 3462

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS

<222> (16)..(3459)

<400> 3

aagggggaga aaagt ttg tct cag caa tcg ata caa aaa gta tta gta gca 51

Leu Ser Gln Gln Ser Ile Gln Lys Val Leu Val Ala

1

5

10

aac agg gga gaa att gca atc cga ata ttc cgg gcg tgt acc gag ttg 99

Asn Arg Gly Glu Ile Ala Ile Arg Ile Phe Arg Ala Cys Thr Glu Leu

15

20

25

aat att cgt aca gtt gcg gtc tat tca aaa gaa gat tcc ggt tcc tac 147

Asn Ile Arg Thr Val Ala Val Tyr Ser Lys Glu Asp Ser Gly Ser Tyr

30

35

40

cat cgg tac aaa gcg gat gaa gca tac ttg gtc ggt gaa ggg aaa aaa 195

His Arg Tyr Lys Ala Asp Glu Ala Tyr Leu Val Gly Glu Gly Lys Lys

45

50

55

60

ccg att gat gct tac ctg gat att gaa ggt atc att gat att gcg aaa 243

Pro Ile Asp Ala Tyr Leu Asp Ile Glu Gly Ile Ile Asp Ile Ala Lys

65

70

75

aga aac aaa gtc gat gca att cat ccg gga tac ggt ttc tta tct gaa 291

Arg Asn Lys Val Asp Ala Ile His Pro Gly Tyr Gly Phe Leu Ser Glu

80

85

90

aat att cat ttt gcg aga cga tgt gaa gaa gaa ggc atc gta ttc ata 339

Asn Ile His Phe Ala Arg Arg Cys Glu Glu Glu Gly Ile Val Phe Ile
 95 100 105
 5 ggg cca aaa tcc gag cat ctc gat atg ttt ggt gac aag gta aaa gcg 387
 Gly Pro Lys Ser Glu His Leu Asp Met Phe Gly Asp Lys Val Lys Ala
 110 115 120
 10 cgt gag cag gca gaa aaa gcg gga atc ccc gtg att ccg gga agc gac 435
 Arg Glu Gln Ala Glu Lys Ala Gly Ile Pro Val Ile Pro Gly Ser Asp
 125 130 135 140
 15 ggt cct gcc gaa acg ctt gaa gcc gtc gaa caa ttt gga caa gct aac 483
 Gly Pro Ala Glu Thr Leu Glu Ala Val Glu Gln Phe Gly Gln Ala Asn
 145 150 155
 20 ggt tat ccg atc atc att aaa gcc tcg ctt ggc ggc ggc ggc cgc ggt 531
 Gly Tyr Pro Ile Ile Ile Lys Ala Ser Leu Gly Gly Gly Gly Arg Gly
 160 165 170
 25 atg cgg att gtc aga tct gaa agt gaa gtt aaa gaa gca tat gag cgt 579
 Met Arg Ile Val Arg Ser Glu Ser Glu Val Lys Glu Ala Tyr Glu Arg
 175 180 185
 30 gct aaa tca gag gcg aaa gca gcc ttt ggc aat gat gaa gtt tat gta 627
 Ala Lys Ser Glu Ala Lys Ala Ala Phe Gly Asn Asp Glu Val Tyr Val
 190 195 200
 35 gaa aaa tta att gag aat ccg aaa cat att gag gtt cag gtc att gga 675
 Glu Lys Leu Ile Glu Asn Pro Lys His Ile Glu Val Gln Val Ile Gly
 205 210 215 220
 40 gac aag cag ggc aat gtc gtc cat ctt ttt gag agg gat tgc tcc gtt 723
 Asp Lys Gln Gly Asn Val Val His Leu Phe Glu Arg Asp Cys Ser Val
 225 230 235
 45 caa aga cgc cat caa aaa gtc att gaa gtg gcg ccg agt gtc tcg ctg 771
 Gln Arg Arg His Gln Lys Val Ile Glu Val Ala Pro Ser Val Ser Leu
 240 245 250
 50 tca cct gaa tta agg gac caa att tgt gag gct gca gtt gcg ctt gcc 819
 Ser Pro Glu Leu Arg Asp Gln Ile Cys Glu Ala Ala Val Ala Leu Ala
 255 260 265
 55 aaa aat gta aac tat ata aat gcg ggg acg gtc gaa ttc ctt gtt gca 867
 Lys Asn Val Asn Tyr Ile Asn Ala Gly Thr Val Glu Phe Leu Val Ala

EP 1 092 776 A1

	270	275	280	
	aac aac gag ttc tac ttt att gaa gta aat cct cgc gta caa gtt gaa			915
5	Asn Asn Glu Phe Tyr Phe Ile Glu Val Asn Pro Arg Val Gln Val Glu			
	285	290	295	300
	cac acg ata aca gaa atg att act ggt gtc gat att gtt caa act cag			963
10	His Thr Ile Thr Glu Met Ile Thr Gly Val Asp Ile Val Gln Thr Gln			
	305	310	315	
	atc ctt gtt gcc caa ggg cac agc ctt cac agc aaa aaa gta aat att			1011
15	Ile Leu Val Ala Gln Gly His Ser Leu His Ser Lys Lys Val Asn Ile			
	320	325	330	
	cct gag caa aag gac att ttt aca atc ggc tat gcc att cag tca cgg			1059
20	Pro Glu Gln Lys Asp Ile Phe Thr Ile Gly Tyr Ala Ile Gln Ser Arg			
	335	340	345	
	gtt acg act gag gat ccg caa aat gat ttc atg cct gat aca gga aaa			1107
25	Val Thr Thr Glu Asp Pro Gln Asn Asp Phe Met Pro Asp Thr Gly Lys			
	350	355	360	
	atc atg gct tac cgc tca ggc ggc ggt ttt ggt gtc cgt ctt gat acc			1155
30	Ile Met Ala Tyr Arg Ser Gly Gly Gly Phe Gly Val Arg Leu Asp Thr			
	365	370	375	380
	gga aac agc ttc cag ggc gcc gtg atc aca cca tac tat gat tca ctt			1203
	Gly Asn Ser Phe Gln Gly Ala Val Ile Thr Pro Tyr Tyr Asp Ser Leu			
	385	390	395	
35	ctc gtt aag ctt tca act tgg gct tta acg ttt gaa cag gca gct gcc			1251
	Leu Val Lys Leu Ser Thr Trp Ala Leu Thr Phe Glu Gln Ala Ala Ala			
	400	405	410	
40	aaa atg gtg cga aac ctt cag gag ttt aga atc aga ggc ata aaa acg			1299
	Lys Met Val Arg Asn Leu Gln Glu Phe Arg Ile Arg Gly Ile Lys Thr			
	415	420	425	
45	aac att ccg ttc ctt gag aac gtt gca aag cat gag aag ttc ctg aca			1347
	Asn Ile Pro Phe Leu Glu Asn Val Ala Lys His Glu Lys Phe Leu Thr			
	430	435	440	
50	ggg caa tat gat aca tct ttc att gat aca acg cct gaa tta ttt aat			1395
	Gly Gln Tyr Asp Thr Ser Phe Ile Asp Thr Thr Pro Glu Leu Phe Asn			
	445	450	455	460

ttc cct aaa caa aaa gac cgc gga acg aaa atg ctc act tac atc ggc 1443
 Phe Pro Lys Gln Lys Asp Arg Gly Thr Lys Met Leu Thr Tyr Ile Gly
 5 465 470 475
 aat gtg aca gtg aac ggc ttc cct gga atc ggg aaa aaa gaa aaa ccg 1491
 Asn Val Thr Val Asn Gly Phe Pro Gly Ile Gly Lys Lys Glu Lys Pro
 10 480 485 490
 gcg ttt gac aaa ccg tta ggc gta aag gta gac gtt gat cag cag cct 1539
 Ala Phe Asp Lys Pro Leu Gly Val Lys Val Asp Val Asp Gln Gln Pro
 15 495 500 505
 gcc aga gga aca aag caa att ctc gat gaa aaa ggt gca gaa ggg ctt 1587
 Ala Arg Gly Thr Lys Gln Ile Leu Asp Glu Lys Gly Ala Glu Gly Leu
 20 510 515 520
 gca aat tgg gtt aag gag cag aaa tct gtc ctt tta act gat acg aca 1635
 Ala Asn Trp Val Lys Glu Gln Lys Ser Val Leu Leu Thr Asp Thr Thr
 25 525 530 535 540
 ttc agg gat gcc cac caa tcg tta ttg gca act aga atc aga tcg cat 1683
 Phe Arg Asp Ala His Gln Ser Leu Leu Ala Thr Arg Ile Arg Ser His
 30 545 550 555
 gat ttg aaa aaa atc gca aat ccg acg gct gcg tta tgg cct gaa cta 1731
 Asp Leu Lys Lys Ile Ala Asn Pro Thr Ala Ala Leu Trp Pro Glu Leu
 35 560 565 570
 ttc agt atg gaa atg tgg gga ggc gcg acc ttc gat gta gcc tac cga 1779
 Phe Ser Met Glu Met Trp Gly Gly Ala Thr Phe Asp Val Ala Tyr Arg
 40 575 580 585
 ttc ctg aaa gaa gat ccg tgg aaa cgt ttg gaa gat ctt cgc aaa gaa 1827
 Phe Leu Lys Glu Asp Pro Trp Lys Arg Leu Glu Asp Leu Arg Lys Glu
 45 590 595 600
 gtg ccg aat acc tta ttc cag atg ttg ctt cgc tca tca aat gcg gtc 1875
 Val Pro Asn Thr Leu Phe Gln Met Leu Leu Arg Ser Ser Asn Ala Val
 50 605 610 615 620
 ggc tat acg aat tat ccg gac aat gtg att aaa gaa ttt gtg aag caa 1923
 Gly Tyr Thr Asn Tyr Pro Asp Asn Val Ile Lys Glu Phe Val Lys Gln
 55 625 630 635
 tca gct caa tcc ggt att gat gtg ttt cgt att ttc gac agc tta aac 1971

EP 1 092 776 A1

Ser Ala Gln Ser Gly Ile Asp Val Phe Arg Ile Phe Asp Ser Leu Asn
640 645 650
5 tgg gta aaa ggg atg acg tta gcc att gat gct gtt agg gat acc ggc 2019
Trp Val Lys Gly Met Thr Leu Ala Ile Asp Ala Val Arg Asp Thr Gly
655 660 665
10 aaa gtg gca gaa gct gcg att tgt tat acg gga gat atc ctt gac aag 2067
Lys Val Ala Glu Ala Ala Ile Cys Tyr Thr Gly Asp Ile Leu Asp Lys
670 675 680
15 aac cgg acg aag tac gac ctt gca tat tat aca tcg atg gcg aag gag 2115
Asn Arg Thr Lys Tyr Asp Leu Ala Tyr Tyr Thr Ser Met Ala Lys Glu
685 690 695 700
20 ctt gag gcg gcc gga gcc cat att ctc ggg att aaa gat atg gca ggc 2163
Leu Glu Ala Ala Gly Ala His Ile Leu Gly Ile Lys Asp Met Ala Gly
705 710 715
25 ctg tta aaa ccg cag gct gca tat gag ctc gtt tct gcg ttg aaa gaa 2211
Leu Leu Lys Pro Gln Ala Ala Tyr Glu Leu Val Ser Ala Leu Lys Glu
720 725 730
30 acg atc gac att ccg gtt cac ctt cat acg cat gat acg agc gga aac 2259
Thr Ile Asp Ile Pro Val His Leu His Thr His Asp Thr Ser Gly Asn
735 740 745
35 ggt att tat atg tat gcg aaa gct gtt gaa gcc ggc gtt gat atc ata 2307
Gly Ile Tyr Met Tyr Ala Lys Ala Val Glu Ala Gly Val Asp Ile Ile
750 755 760
40 gac gtg gcg gtc agc tca atg gcg gga tta acg tca cag cct agc gcg 2355
Asp Val Ala Val Ser Ser Met Ala Gly Leu Thr Ser Gln Pro Ser Ala
765 770 775 780
45 agc gga ttt tat cat gcg atg gaa ggc aac gac cgc cgt ccg gaa atg 2403
Ser Gly Phe Tyr His Ala Met Glu Gly Asn Asp Arg Arg Pro Glu Met
785 790 795
50 aat gtc caa ggc gtt gaa ttg ctg tcc caa tat tgg gag tgc gtg cgt 2451
Asn Val Gln Gly Val Glu Leu Leu Ser Gln Tyr Trp Glu Ser Val Arg
800 805 810
55 aaa tat tat agt gaa ttt gaa agc gga atg aag tct ccg cat act gaa 2499
Lys Tyr Tyr Ser Glu Phe Glu Ser Gly Met Lys Ser Pro His Thr Glu

EP 1 092 776 A1

	815	820	825	
	att tat gaa cac gaa atg cca ggg ggc caa tac agc aac ctg cag cag	2547		
5	Ile Tyr Glu His Glu Met Pro Gly Gly Gln Tyr Ser Asn Leu Gln Gln			
	830	835	840	
	caa gcc aag gga gta ggc ctt ggc gac cgc tgg aac gaa gtc aag gaa	2595		
10	Gln Ala Lys Gly Val Gly Leu Gly Asp Arg Trp Asn Glu Val Lys Glu			
	845	850	855	860
	atg tac aga cgc gtg aac gat atg ttc ggt gac atc gtc aag gta acg	2643		
15	Met Tyr Arg Arg Val Asn Asp Met Phe Gly Asp Ile Val Lys Val Thr			
	865	870	875	
	cct tcc tca aaa gta gtc gga gat atg gca ctc tac atg gtg caa aac	2691		
20	Pro Ser Ser Lys Val Val Gly Asp Met Ala Leu Tyr Met Val Gln Asn			
	880	885	890	
	aat ctg act gaa aaa gac gtt tac gaa aaa ggt gaa tct tta gat ttc	2739		
	Asn Leu Thr Glu Lys Asp Val Tyr Glu Lys Gly Glu Ser Leu Asp Phe			
25	895	900	905	
	cct gat tct gtc gtg gag ctt ttt aaa gga aat atc ggc cag cct cat	2787		
	Pro Asp Ser Val Val Glu Leu Phe Lys Gly Asn Ile Gly Gln Pro His			
30	910	915	920	
	ggc gga ttc cca gaa aaa ctg caa aag ctg atc tta aaa ggg cag gag	2835		
	Gly Gly Phe Pro Glu Lys Leu Gln Lys Leu Ile Leu Lys Gly Gln Glu			
	925	930	935	940
35	ccg att aca gtc aga ccg ggc gaa ctg ctt gag ccg gtg tca ttt gaa	2883		
	Pro Ile Thr Val Arg Pro Gly Glu Leu Leu Glu Pro Val Ser Phe Glu			
	945	950	955	
40	gcg atc aaa cag gaa ttt aaa gag cag cat aac ttg gaa att tca gat	2931		
	Ala Ile Lys Gln Glu Phe Lys Glu Gln His Asn Leu Glu Ile Ser Asp			
	960	965	970	
45	cag gat gct gtg gca tat gcc ctt tat cct aaa gtc ttc act gat tat	2979		
	Gln Asp Ala Val Ala Tyr Ala Leu Tyr Pro Lys Val Phe Thr Asp Tyr			
	975	980	985	
50	gtg aaa acg aca gaa agc tat gga gac atc tcg gta tta gat aca ccg	3027		
	Val Lys Thr Thr Glu Ser Tyr Gly Asp Ile Ser Val Leu Asp Thr Pro			
	990	995	1000	

EP 1 092 776 A1

5
10
15
20
25
30
35
40
45

aca ttc ttc tac ggt atg aca tta ggt gaa gag ata gaa gtt gaa att 3075
Thr Phe Phe Tyr Gly Met Thr Leu Gly Glu Glu Ile Glu Val Glu Ile
1005 1010 1015 1020
gag cgc ggc aaa acg ctg atc gtt aag ctg att tca atc ggt gag cct 3123
Glu Arg Gly Lys Thr Leu Ile Val Lys Leu Ile Ser Ile Gly Glu Pro
1025 1030 1035
cag cct gat gcc acc cgc gtc gtt tat ttc gaa ctc aac ggg cag ccg 3171
Gln Pro Asp Ala Thr Arg Val Val Tyr Phe Glu Leu Asn Gly Gln Pro
1040 1045 1050
cgt gaa gta gtc att aaa gat gaa agc att aag tct tcc gtt cag gaa 3219
Arg Glu Val Val Ile Lys Asp Glu Ser Ile Lys Ser Ser Val Gln Glu
1055 1060 1065
agg ctg aaa gca gac cgg aca aat cca agc cac atc gca gct tcc atg 3267
Arg Leu Lys Ala Asp Arg Thr Asn Pro Ser His Ile Ala Ala Ser Met
1070 1075 1080
cct gga aca gtt att aag gta ttg gct gaa gca ggc aca aaa gtc aat 3315
Pro Gly Thr Val Ile Lys Val Leu Ala Glu Ala Gly Thr Lys Val Asn
1085 1090 1095 1100
aaa ggt gat cat ttg atg att aat gaa gcg atg aaa atg gaa aca acg 3363
Lys Gly Asp His Leu Met Ile Asn Glu Ala Met Lys Met Glu Thr Thr
1105 1110 1115
gtt cag gcg cct ttc tca gga aca atc aag cag gtt cat gtg aaa aat 3411
Val Gln Ala Pro Phe Ser Gly Thr Ile Lys Gln Val His Val Lys Asn
1120 1125 1130
ggt gag ccg atc caa acg gga gat ctg ctc ctt gaa att gaa aaa gca 3459
Gly Glu Pro Ile Gln Thr Gly Asp Leu Leu Leu Glu Ile Glu Lys Ala
1135 1140 1145
taa 3462

<210> 4

<211> 1148

<212> PRT

<213> Bacillus subtilis

<400> 4

5 Leu Ser Gln Gln Ser Ile Gln Lys Val Leu Val Ala Asn Arg Gly Glu
 1 5 10 15
 Ile Ala Ile Arg Ile Phe Arg Ala Cys Thr Glu Leu Asn Ile Arg Thr
 20 25 30
 10 Val Ala Val Tyr Ser Lys Glu Asp Ser Gly Ser Tyr His Arg Tyr Lys
 35 40 45
 Ala Asp Glu Ala Tyr Leu Val Gly Glu Gly Lys Lys Pro Ile Asp Ala
 50 55 60
 15 Tyr Leu Asp Ile Glu Gly Ile Ile Asp Ile Ala Lys Arg Asn Lys Val
 65 70 75 80
 20 Asp Ala Ile His Pro Gly Tyr Gly Phe Leu Ser Glu Asn Ile His Phe
 85 90 95
 Ala Arg Arg Cys Glu Glu Glu Gly Ile Val Phe Ile Gly Pro Lys Ser
 100 105 110
 25 Glu His Leu Asp Met Phe Gly Asp Lys Val Lys Ala Arg Glu Gln Ala
 115 120 125
 Glu Lys Ala Gly Ile Pro Val Ile Pro Gly Ser Asp Gly Pro Ala Glu
 130 135 140
 30 Thr Leu Glu Ala Val Glu Gln Phe Gly Gln Ala Asn Gly Tyr Pro Ile
 145 150 155 160
 Ile Ile Lys Ala Ser Leu Gly Gly Gly Gly Arg Gly Met Arg Ile Val
 35 165 170 175
 Arg Ser Glu Ser Glu Val Lys Glu Ala Tyr Glu Arg Ala Lys Ser Glu
 180 185 190
 40 Ala Lys Ala Ala Phe Gly Asn Asp Glu Val Tyr Val Glu Lys Leu Ile
 195 200 205
 Glu Asn Pro Lys His Ile Glu Val Gln Val Ile Gly Asp Lys Gln Gly
 210 215 220
 45 Asn Val Val His Leu Phe Glu Arg Asp Cys Ser Val Gln Arg Arg His
 225 230 235 240
 50 Gln Lys Val Ile Glu Val Ala Pro Ser Val Ser Leu Ser Pro Glu Leu
 245 250 255
 Arg Asp Gln Ile Cys Glu Ala Ala Val Ala Leu Ala Lys Asn Val Asn

EP 1 092 776 A1

260 265 270
 Tyr Ile Asn Ala Gly Thr Val Glu Phe Leu Val Ala Asn Asn Glu Phe
 5 275 280 285
 Tyr Phe Ile Glu Val Asn Pro Arg Val Gln Val Glu His Thr Ile Thr
 290 295 300
 10 Glu Met Ile Thr Gly Val Asp Ile Val Gln Thr Gln Ile Leu Val Ala
 305 310 315 320
 Gln Gly His Ser Leu His Ser Lys Lys Val Asn Ile Pro Glu Gln Lys
 325 330 335
 15 Asp Ile Phe Thr Ile Gly Tyr Ala Ile Gln Ser Arg Val Thr Thr Glu
 340 345 350
 Asp Pro Gln Asn Asp Phe Met Pro Asp Thr Gly Lys Ile Met Ala Tyr
 20 355 360 365
 Arg Ser Gly Gly Gly Phe Gly Val Arg Leu Asp Thr Gly Asn Ser Phe
 370 375 380
 25 Gln Gly Ala Val Ile Thr Pro Tyr Tyr Asp Ser Leu Leu Val Lys Leu
 385 390 395 400
 Ser Thr Trp Ala Leu Thr Phe Glu Gln Ala Ala Ala Lys Met Val Arg
 405 410 415
 30 Asn Leu Gln Glu Phe Arg Ile Arg Gly Ile Lys Thr Asn Ile Pro Phe
 420 425 430
 Leu Glu Asn Val Ala Lys His Glu Lys Phe Leu Thr Gly Gln Tyr Asp
 35 435 440 445
 Thr Ser Phe Ile Asp Thr Thr Pro Glu Leu Phe Asn Phe Pro Lys Gln
 450 455 460
 40 Lys Asp Arg Gly Thr Lys Met Leu Thr Tyr Ile Gly Asn Val Thr Val
 465 470 475 480
 Asn Gly Phe Pro Gly Ile Gly Lys Lys Glu Lys Pro Ala Phe Asp Lys
 485 490 495
 45 Pro Leu Gly Val Lys Val Asp Val Asp Gln Gln Pro Ala Arg Gly Thr
 500 505 510
 Lys Gln Ile Leu Asp Glu Lys Gly Ala Glu Gly Leu Ala Asn Trp Val
 50 515 520 525
 Lys Glu Gln Lys Ser Val Leu Leu Thr Asp Thr Thr Phe Arg Asp Ala

EP 1 092 776 A1

	530	535	540
	His Gln Ser Leu Leu Ala Thr Arg Ile Arg Ser His Asp Leu Lys Lys		
5	545	550	555 560
	Ile Ala Asn Pro Thr Ala Ala Leu Trp Pro Glu Leu Phe Ser Met Glu		
	565	570	575
10	Met Trp Gly Gly Ala Thr Phe Asp Val Ala Tyr Arg Phe Leu Lys Glu		
	580	585	590
	Asp Pro Trp Lys Arg Leu Glu Asp Leu Arg Lys Glu Val Pro Asn Thr		
15	595	600	605
	Leu Phe Gln Met Leu Leu Arg Ser Ser Asn Ala Val Gly Tyr Thr Asn		
	610	615	620
	Tyr Pro Asp Asn Val Ile Lys Glu Phe Val Lys Gln Ser Ala Gln Ser		
20	625	630	635 640
	Gly Ile Asp Val Phe Arg Ile Phe Asp Ser Leu Asn Trp Val Lys Gly		
	645	650	655
25	Met Thr Leu Ala Ile Asp Ala Val Arg Asp Thr Gly Lys Val Ala Glu		
	660	665	670
	Ala Ala Ile Cys Tyr Thr Gly Asp Ile Leu Asp Lys Asn Arg Thr Lys		
30	675	680	685
	Tyr Asp Leu Ala Tyr Tyr Thr Ser Met Ala Lys Glu Leu Glu Ala Ala		
	690	695	700
	Gly Ala His Ile Leu Gly Ile Lys Asp Met Ala Gly Leu Leu Lys Pro		
35	705	710	715 720
	Gln Ala Ala Tyr Glu Leu Val Ser Ala Leu Lys Glu Thr Ile Asp Ile		
	725	730	735
40	Pro Val His Leu His Thr His Asp Thr Ser Gly Asn Gly Ile Tyr Met		
	740	745	750
	Tyr Ala Lys Ala Val Glu Ala Gly Val Asp Ile Ile Asp Val Ala Val		
45	755	760	765
	Ser Ser Met Ala Gly Leu Thr Ser Gln Pro Ser Ala Ser Gly Phe Tyr		
	770	775	780
	His Ala Met Glu Gly Asn Asp Arg Arg Pro Glu Met Asn Val Gln Gly		
50	785	790	795 800
	Val Glu Leu Leu Ser Gln Tyr Trp Glu Ser Val Arg Lys Tyr Tyr Ser		

55

EP 1 092 776 A1

	805	810	815
	Glu Phe Glu Ser Gly Met Lys Ser Pro His Thr Glu Ile Tyr Glu His		
5	820	825	830
	Glu Met Pro Gly Gly Gln Tyr Ser Asn Leu Gln Gln Gln Ala Lys Gly		
	835	840	845
10	Val Gly Leu Gly Asp Arg Trp Asn Glu Val Lys Glu Met Tyr Arg Arg		
	850	855	860
	Val Asn Asp Met Phe Gly Asp Ile Val Lys Val Thr Pro Ser Ser Lys		
15	865	870	875
	Val Val Gly Asp Met Ala Leu Tyr Met Val Gln Asn Asn Leu Thr Glu		
	885	890	895
	Lys Asp Val Tyr Glu Lys Gly Glu Ser Leu Asp Phe Pro Asp Ser Val		
20	900	905	910
	Val Glu Leu Phe Lys Gly Asn Ile Gly Gln Pro His Gly Gly Phe Pro		
	915	920	925
25	Glu Lys Leu Gln Lys Leu Ile Leu Lys Gly Gln Glu Pro Ile Thr Val		
	930	935	940
	Arg Pro Gly Glu Leu Leu Glu Pro Val Ser Phe Glu Ala Ile Lys Gln		
30	945	950	955
	Glu Phe Lys Glu Gln His Asn Leu Glu Ile Ser Asp Gln Asp Ala Val		
	965	970	975
	Ala Tyr Ala Leu Tyr Pro Lys Val Phe Thr Asp Tyr Val Lys Thr Thr		
35	980	985	990
	Glu Ser Tyr Gly Asp Ile Ser Val Leu Asp Thr Pro Thr Phe Phe Tyr		
	995	1000	1005
40	Gly Met Thr Leu Gly Glu Glu Ile Glu Val Glu Ile Glu Arg Gly Lys		
	1010	1015	1020
	Thr Leu Ile Val Lys Leu Ile Ser Ile Gly Glu Pro Gln Pro Asp Ala		
45	1025	1030	1035
	Thr Arg Val Val Tyr Phe Glu Leu Asn Gly Gln Pro Arg Glu Val Val		
	1045	1050	1055
	Ile Lys Asp Glu Ser Ile Lys Ser Ser Val Gln Glu Arg Leu Lys Ala		
50	1060	1065	1070
	Asp Arg Thr Asn Pro Ser His Ile Ala Ala Ser Met Pro Gly Thr Val		

1075 1080 1085
 Ile Lys Val Leu Ala Glu Ala Gly Thr Lys Val Asn Lys Gly Asp His
 5 1090 1095 1100
 Leu Met Ile Asn Glu Ala Met Lys Met Glu Thr Thr Val Gln Ala Pro
 105 1110 1115 1120
 10 Phe Ser Gly Thr Ile Lys Gln Val His Val Lys Asn Gly Glu Pro Ile
 1125 1130 1135
 Gln Thr Gly Asp Leu Leu Leu Glu Ile Glu Lys Ala
 15 1140 1145

Claims

1. A bacterium belonging to the genus *Escherichia* which contains a gene coding for an enzyme having pyruvate carboxylase activity and which has L-amino acid productivity.
2. The bacterium according to claim 1, wherein the gene coding for an enzyme having pyruvate carboxylase activity is pyruvate carboxylase derived from a bacterium belonging to the genus *Bacillus*.
3. The bacterium according to claim 1 or 2, wherein said gene has the nucleotide sequence as shown in SEQ ID NO 3.
4. The bacterium according to any of the preceding claims, wherein said enzyme has an amino acid sequence as shown in SEQ ID NO 4.
5. The bacterium according to any of the preceding claims which is transformed with a vector containing said gene.
6. The bacterium according to any of the preceding claims, wherein the gene is introduced into said bacterium in a low copy number.
7. The bacterium according to any of the preceding claims, wherein the L-amino acid is selected from the group consisting of L-threonine, L-lysine, L-glutamic acid, L-homoserine, L-methionine, L-arginine, L-proline and L-isoleucine.
8. The bacterium according to any of the preceding claims which produces a higher amount of the L-amino acid than the bacterium not containing said gene.
9. The bacterium according to any of the preceding claims which produces L-threonine in a concentration which is at least 100 % higher than the concentration of L-threonine produced by a bacterium not containing said gene.
10. The bacterium according to any of the preceding claims which produces L-glutamic acid in a concentration which is at least 50 % higher than the concentration of L-glutamic acid produced by a bacterium not containing said gene.
11. The bacterium according to any of the preceding claims which produces L-homoserine in a concentration which is at least 25 % higher than the concentration of L-homoserine produced by a bacterium not containing said gene.
12. The bacterium having the accession number VKPM B-7821.
13. The bacterium having the accession number VKPM B-7822.

EP 1 092 776 A1

14. A method for producing an L-amino acid comprising cultivating the bacterium according to any of the claims 1 to 13 in a medium, producing and accumulating the L-amino acid in the medium.

5

10

15

20

25

30

35


40

45

50

55

PCR with oligonucleotides: 5'-GATATAAGGGGACTTCAGAG and 5'-GGCGCTTTATGCGTTTCAATC
 ↓ and *B. subtilis* 168 chromosomal DNA



Transformation of *B. subtilis* 168 trpC2; and selection of CmR-clone of integrant
B. subtilis trpC2 xyz(269 b.p.):pBRPYCA11CmR3;




Fig. 1

B. subtilis trpC2 pycA::KmR (phenotype Asp(-) KmR)
 ↓
 Transduction with phages E40 lysate prepared on
B. subtilis recE::tet (TcR) strain;
 Selection of TcR clones of
B. subtilis trpC2 pycA::KmR recE::TcR
 (phenotype Asp(-) Trp(-) KmR TcR)
 This strain was used as the recipient for the cloning of pycA gene.

Chromosomal DNA of *B. subtilis* 168 trpC2 was partially digested with EcoRI.





Fig.2



25



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 12 1763

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCL.7)
Y	WO 99 18228 A (PETERS WENDISCH PETRA ;EIKMANN BERND (DE); SAHM HERMANN (DE); KER) 15 April 1999 (1999-04-15) * page 1, line 24-30 * * page 3, line 7 - page 5, line 12 * * examples 4-6 * * claims 1,4,6,32,34 * ---	1-14	C12N15/52 C12N9/00 C12P13/06 C12P13/08 C12P13/14 C12R1/19 //(C12P13/06, C12R1:19), (C12P13/08, C12R1:19), (C12P13/14, C12R1:19)
Y	PETERS-WENDISCH ET AL: "Pyruvate carboxylase as an anaplerotic enzyme in Corynebacterium glutamicum" MICROBIOLOGY, vol. 143, no. PART 04, April 1997 (1997-04), pages 1095-1103, XP002110209 * abstract * * page 1095, right-hand column, line 6 - page 1096, left-hand column, line 21 * * page 1100, right-hand column, line 28-47 * ---	1-14	
Y	DATABASE EMBL 'Online! EMBL; ID BSUB0008, AC Z99111, 20 November 1997 (1997-11-20) KUNST F ET AL.: "Bacillus subtilis complete genome (section 8 of 21); from 1394791 to 1603020" XP002160795 * page 36, line 28-59 * * page 85 - page 86 * --- -/--	1-14	TECHNICAL FIELDS SEARCHED (InCL.7) C12N C12P C12R
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 20 February 2001	Examiner van de Kamp, M
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1500 03.02 (P04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 12 1763

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	KOFFAS ET AL: "Sequence of the Corynebacterium glutamicum pyruvate carboxylase gene" APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, vol. 50, no. 3, September 1998 (1998-09), pages 346-352, XP002110411 * abstract * * page 346, right-hand column, line 11 - page 347, left-hand column, line 23 *	1-8,14	
A	FR 2 738 014 A (AJINOMOTO KK) 28 February 1997 (1997-02-28) * examples 3-5 *	1,14	
D,A	US 4 278 765 A (DEBABOV VLADIMIR G ET AL) 14 July 1981 (1981-07-14) * example 1 *	12	
A	KRAMER R: "Genetic and physiological approaches for the production of amino acids" JOURNAL OF BIOTECHNOLOGY, vol. 45, no. 1, 12 February 1996 (1996-02-12), pages 1-21, XP004036833 * abstract; figures 2,3 *	14	
P,X	WO 99 53035 A (ALTMAN ELLIOT ;GOKARN RAVI R (US); EITEMAN MARK A (US); UNIV GEORG) 21 October 1999 (1999-10-21) * example 4 * * claims 1-3,6,18,42,45,46,49 *	1,5,7-9,14	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 20 February 2001	Examiner van de Kamp, M
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons 8 : member of the same patent family, corresponding document	

EPO FORM 1503 03.82 [P4/C01]

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 12 1763

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20-02-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9918228 A	15-04-1999	DE 19831609 A	15-04-1999
		AU 1148299 A	27-04-1999
		BR 9813021 A	15-08-2000
		CN 1275167 T	29-11-2000
		EP 1015621 A	05-07-2000
FR 2738014 A	28-02-1997	BR 9610016 A	06-07-1999
		CN 1193343 A	16-09-1998
		DE 19681532 T	03-12-1998
		WO 9708294 A	06-03-1997
		US 6110714 A	29-08-2000
US 4278765 A	14-07-1981	SU 875663 A	15-09-1982
		HU 190999 B	28-12-1986
WO 9953035 A	21-10-1999	AU 3555999 A	01-11-1999
		EP 1073722 A	07-02-2001

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82